



Identification of novel susceptibility genes in ozone-induced inflammation in mice

A.K. Bauer*, E.L. Travis[#], S.S. Malhotra[#], E.A. Rondini*, C. Walker[†], H-Y. Cho[†], S. Trivedi[†], W. Gladwell[†], S. Reddy⁺ and S.R. Kleeberger[†]

ABSTRACT: Ozone (O₃) remains a prevalent air pollutant and public health concern. *Inf2* is a significant quantitative trait locus on murine chromosome 17 that contributes to susceptibility to O₃-induced infiltration of polymorphonuclear leukocytes (PMNs) into the lung, but the mechanisms of susceptibility remain unclear. The study objectives were to confirm and restrict *Inf2*, and to identify and test novel candidate susceptibility gene(s).

Congenic strains of mice that contained overlapping regions of *Inf2* and their controls, and mice deficient in either major histocompatibility complex (MHC) class II genes or the *Tnf* cluster, were exposed to air or O₃. Lung inflammation and gene expression were assessed.

Inf2 was restricted from 16.42 Mbp to 0.96 Mbp, and bioinformatic analysis identified MHC class II, the *Tnf* cluster and other genes in this region that contain potentially informative single nucleotide polymorphisms between the susceptible and resistant mice. Furthermore, O₃-induced inflammation was significantly reduced in mice deficient in MHC class II genes or the *Tnf* cluster genes, compared with wild-type controls. Gene expression differences were also observed in MHC class II and *Tnf* cluster genes.

This integrative genetic analysis of *Inf2* led to identification of novel O₃ susceptibility genes that may provide important, new therapeutic targets in susceptible individuals.

KEYWORDS: Inflammation, lymphotoxin α , major histocompatibility complex, mouse, susceptibility, tumour necrosis factor

The pollutant ozone (O₃) is a highly toxic principal oxidant found in urban environments throughout industrialised cities worldwide. O₃ exposure has been associated with many adverse health effects, such as exacerbation of asthma [1, 2]. Identification of susceptibility genes involved in O₃-induced pulmonary injury may provide critical information for future risk assessment as well as general international health policies. In 2006, an estimated one-third of US individuals were at an increased risk of adverse effects caused by O₃ and 131 million US residents resided in regions that either approached or exceeded the National Ambient Air Quality Standard of 0.08 ppm O₃ [3].

Significant intersubject differences in pulmonary function and inflammatory responses to O₃ suggest that genetic background contributes to O₃ susceptibility in humans and rodents [4–6]. Furthermore, studies in human subjects have suggested that polymorphisms in oxidant defence genes, such as glutathione S-transferase

M1, and quinone metabolism genes, such as NADPH quinone oxidoreductase 1, associate with differential responsiveness to O₃ [7, 8]. Activating polymorphisms in inflammatory genes, such as tumour necrosis factor alpha (TNF), also enhance susceptibility to O₃ and asthma [9]. A genome-wide linkage analysis of O₃-induced influx of polymorphonuclear leukocytes (PMNs) in an intercross cohort (B6C3F₂) derived from susceptible C57BL/6 (B6) and resistant C3H/HeJ (C3) progenitor mouse strains identified a significant susceptibility quantitative trait locus (QTL) on chromosome 17 (inflammation 2; *Inf2*) [5]. *Inf2* (33.73–50.15 Mbp; D17Mit16–D17Mit10) contains the *H-2* locus, including major histocompatibility complex (MHC) genes and non-MHC genes, such as the pro-inflammatory cytokine *Tnf*. Pre-treatment of B6 mice with a monoclonal antibody to TNF- α and deletion of TNF- α receptors 1 and 2 [5, 10] significantly attenuated the inflammatory response to O₃ relative to control mice, supporting the importance of *Tnf* as a candidate gene in this model.

AFFILIATIONS

*Dept of Pathobiology and Diagnostic Investigation, Michigan State University, East Lansing, MI, [#]University of Texas MD Anderson Cancer Center, Houston, TX, [†]Laboratory of Respiratory Biology, NIEHS, Research Triangle Park, NC, and ⁺Johns Hopkins University, Baltimore, MD, USA.

CORRESPONDENCE

A.K. Bauer
Dept of Pathobiology and Diagnostic Investigation
Center for Integrative Toxicology
B43 Food Safety and Toxicology Bldg
Michigan State University
East Lansing
MI 48824
USA
E-mail: akbauer@msu.edu

Received:

Sept 13 2009

Accepted after revision:

Nov 22 2009

First published online:

Dec 23 2009

This article has supplementary material available at www.erj.ersjournals.com

However, these studies did not conclusively identify *Tnf* as the susceptibility gene in *Inf2*. In addition, *Inf2* is located in the most gene dense, polymorphic region of the entire mouse genome, thus making single candidate gene identification difficult [11].

In the current study we sought to confirm the importance of *Inf2* in O₃-induced lung inflammation, to reduce *Inf2*, and to identify candidate susceptibility genes. Using an integrative genomics approach, we utilised congenic mouse strains to limit *Inf2* from 16.42 Mbp to 0.96 Mbp, and bioinformatic analysis which identified MHC class II genes and the entire *Tnf* cluster as candidate susceptibility loci. Functional analyses of these genes confirmed novel roles for modulation of the inflammatory response to O₃ exposure.

MATERIALS AND METHODS

Mouse strains and O₃ exposure

The following male (6–8 week) congenic mice were used: B10.A-H2^{h2}/(2R)SgSnJ (2R); B10.A-H2^{h4}/(4R)SgDvEg (4R); B10.A-H2ⁱ⁵-H2-T18^a/(5R)SgSnJ (5R); and C3.SW-H2b/SnJ (C3H-H2^b). The congenic region and haplotype for each of the strains are shown in table 1 and figure 1. Control strains for the congenic mice were C57BL/10SnJ (B10), A/WySnJ (A), and C3H/HeSnJ (HeSnJ). The location of the congenic region of the 2R, 4R, and 5R mice with respect to the B10 and A/Wy background strains was identified by genotyping (S.S. Malhotra and E.L. Travis; data not shown). H2^k haplotypes are O₃-resistant and the H2^b haplotype is O₃-susceptible [5]. Additional strains used for the candidate gene studies were B6.129-H2^{dIAbl-Ea}/J (H2^{dIAbl-Ea}; B6 background strain), B6.129S2-Lta^{tm1Dch}/J (Lta^{-/-}; B6 background), B6.129P2-Ltb/Tnf/Lta^{tm1Dck}/J (Ltb/Tnf/Lta^{-/-} mice, B6 background), C57BL/6J (B6) mice, and C3H/HeJ (C3) mice. B6 are O₃-susceptible and are H2^b, while C3H/HeJ (C3) mice are O₃-resistant and are H2^k. All mice were purchased from Jackson Laboratory (Bar Harbor, ME, USA). All animal use was conducted in facilities accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care and approved by

the National Institute of Environmental Health Sciences Animal Care and Use Committee and follows the Helsinki convention for the use and care of animals. For air and O₃ exposures, mice were caged in a humidity- and temperature-controlled room and provided water and pelleted open-formula rodent diet NIH-07 (Zeigler Brothers, Gardners, PA, USA) *ad libitum* (see O₃ exposure procedures in the online supplementary material).

Bronchoalveolar lavage fluid analysis

The procedures used for these techniques have been described previously for right lung lavages and inflammatory cell analysis [5, 10, 14].

Total RNA isolation and real-time quantitative RT-PCR

Mice were sacrificed immediately after air or O₃ exposure, and lungs of each animal were snap-frozen in liquid nitrogen. Total RNA isolation, reverse transcription into cDNA and PCR reaction procedures are described in the online supplementary materials.

Statistics

Data are expressed as group mean ± SEM. Two-way ANOVA was used to evaluate the effects of exposure (air *versus* O₃) and strain (B10, 2R, 4R, 5R, A, HeSnJ, C3H-H2^b, H2^{dIAbl-Ea}+/+, H2^{dIAbl-Ea}+/+, Lta^{-/-}, Lta^{+/+}, Lta/Tnf/Ltb^{-/-}, Lta/Tnf/Ltb^{+/+}, B6, C3) on bronchoalveolar lavage (BAL) phenotypes and mRNA expression. The figure legends contain the number of mice used per experiment. Student–Newman–Keuls test was used for *a posteriori* comparisons of means. All analyses were performed using a commercial statistical analysis package (SigmaStat; Jandel Scientific Software, San Rafael, CA, USA). Statistical significance was accepted at *p* < 0.05.

RESULTS

Restriction of *Inf2*

Responses to O₃ in congenic mice

To confirm the significance of *Inf2*, C3H-H2^b mice which contain *Inf2* from a susceptible mouse (H2^b) on an O₃-resistant

TABLE 1 Nomenclature and characteristics of the congenic and control strains used for investigation

Strain name	Abbreviation	Congenic region		H2 haplotype [#]
		Mbp	cM	
Congenic strains				
B10.A-H2 ^{h2} /(2R)SgSnJ	2R	33.74–35.34	18.20–19.06	H2 ^{h2}
B10.A-H2 ^{h4} /(4R)SgDvEgJ	4R	29.09–34.38	16.30–18.64	H2 ^{h4}
B10.A-H2 ⁱ⁵ -H2-T18 ^a /(5R)SgSnJ	5R	34.45–36.35	18.70–19.14	H2 ⁱ⁵
C3.SW-H2 ^b /SnJ	C3H-H2 ^b	33.74–43.80	18.20–23.30	H2 ^{bc}
Control strains				
C57BL/10SnJ [*]	B10			H2 ^b
A/WySnJ ⁺	A			H2 ^a
C3H/HeSnJ ⁵	HeSnJ			H2 ^k

Mbp and cM were determined using the Mouse Genome Informatics website [12] and genotyped by S.S. Malhotra and E.L. Travis; data not shown. [#]: major histocompatibility complex (MHC) H2 haplotypes according to MHC H2 Haplotype Appendix 2 [13]; ^{*}: background strain for 2R, 4R and 5R mice; ⁺: donor strain for 2R, 4R and 5R mice; ⁵: background strain for C3H-H2^b mice.




FIGURE 1. Schematic of the chromosome 17 congenic segments for 2R, 4R, 5R and C3H-H2^b mice that were used to reduce inflammation 2 (*Inf2*). The congenic segments are shown with respect to *Inf2* (on left). The open rectangle represents the reduced *Inf2*. Mouse genes and human homologues and their chromosomal locations within the reduced *Inf2* are shown on the right. While most of the genes are in the same order in both species, some are not. These locations were identified using the Mouse Genome Informatics website [12] and National Centre for Biotechnology Information. The grey shaded genes are major histocompatibility complex (MHC) class II genes and *Tnf* cluster genes identified as candidate genes using proof of concept experiments (see also figures 5 and 6). *Ager*: advanced glycosylation end product-specific receptor; *Agpat1*: 1-acylglycerol-3-phosphate O-acyltransferase 1; *Aif1*: allograft inflammatory factor 1; *Atf6b*: activating transcription factor 6 beta; *Bat5*: HLA-B associated transcript 5; *Btnl2*: butyrophilin-like 2; *Btnl1*: butyrophilin-like 1; *C2*: complement component 2; *C4a*: complement component 4A; *C4b*: complement component 4B; *Cfb*: complement factor B; *CREBL1*: cAMP responsive element binding protein-like 1; *Cyp21a1*: cytochrome P450, family 21, subfamily a, polypeptide 1; *Cyp21a2-ps*: cytochrome P450, family 21, subfamily a, polypeptide 2 pseudogene; *Ddah*: dimethylarginine dimethylaminohydrolase 2; *Egfl8*: EGF-like domain 8; *Ehmt2*: euchromatic histone-lysine N-methyltransferase 2; *Fkbp1*: FK506 binding protein-like; *Gpsm3*: G-protein signalling modulator 3; *H2-Ab1*: class II antigen A, beta 1; *H2-Aa*: class II antigen A, alpha; *H2-Eb1*: class II antigen E beta; *H2-Eb2*: class II antigen E beta 2; *H2-Ea*: class II antigen E alpha; *H2-ob*: O region beta locus; *Hspa1b*: heat shock 70 kDa protein 1B; *Hspa1a*: heat shock 70 kDa protein 1A; *Lsm2*: LSM2 homolog; *Lta*: lymphotoxin a; *Ltb*: lymphotoxin b; *Ly6g6c*: lymphocyte antigen 6 complex, locus G6C; *Ly6g5c*: lymphocyte antigen 6 complex, locus G5C; *Msh5*: mutS homologue 5; *Neu1*: neuraminidase 1; *Notch4*: notch gene homologue 4; *Pbx2*: pre B-cell leukaemia transcription factor 2; *Ppt2*: palmitoyl-protein thioesterase 2; *Prrt1*: proline-rich transmembrane protein 1; *Psm8*: proteosome subunit, beta type 8; *Rdbp*: RD RNA-binding protein; *Rnf5*: ring finger protein 5; *Slc44a4*: solute carrier family 44, member 4; *Stk19*: serine/threonine kinase 19; *Tap-1*: transporter 1, ATP-binding cassette, sub-family B (MDR/TAP); *Tap-2*: transporter 2, ATP-binding cassette, sub-family B (MDR/TAP); *Tnf*: tumour necrosis factor; *Tnxb*: tenascin B; *Vars*: valyl-tRNA synthetase; *Zbtb12*: zinc finger and BTB domain containing 12. No human homolog exists for *Btnl1*.

H2^K background (HeSnJ; fig. 1) were exposed to air and O₃. Relative to HeSnJ controls, significant increases in mean BAL protein concentration and numbers of macrophages, PMNs and epithelial cells were found in C3H-H2^b mice after 48 and 72 h exposure to O₃ (fig. 2); no strain differences were observed in any parameter in the air-exposed mice. Results thus confirmed the importance of the region of chromosome 17 encompassed by *Inf2* in susceptibility to O₃-induced inflammation.

To reduce the length of *Inf2*, O₃-induced inflammatory responses in congenic 2R, 4R and 5R mice were compared with those of similarly exposed O₃-susceptible background B10 and O₃-resistant donor A strains. Each congenic strain contains a different region of the *Inf2* locus from the A strain mouse on a B10 background (table 1). O₃ caused significant increases in mean numbers of macrophages and PMNs in B10 but not A mice compared with respective air-exposed animals, and numbers of macrophages and PMNs were greater in B10 mice compared with A mice after O₃ (fig. 3). The mean numbers of PMNs and macrophages in 2R and 5R mice were not different from A strain mice after 48 and 72 h O₃; however, numbers of PMNs and macrophages in 2R and 5R mice were significantly lower compared to B10 mice (fig. 3). In contrast, the numbers of BAL fluid (BALF) macrophages and PMNs from O₃-exposed 4R mice were not significantly different from those from O₃-exposed B10 mice, and were greater than those from A, 2R and 5R mice (fig. 3). The mean BALF protein concentrations were significantly reduced in 2R, 4R and 5R mice compared to B10 mice and significantly greater than A mice after 48 and 72 h O₃. Comparison of the mean numbers of macrophages and PMNs between congenic strains with respective congenic regions thus suggested that the 34.38–35.34 Mbp (18.64–19.06 cM) region of chromosome 17 accounted for a major portion of the inflammatory response to O₃ in these mice (fig. 1). No differences in mean numbers of BALF epithelial cells were found between the congenic, A or B10 strains after exposure to O₃ (fig. 3c).

Candidate gene analysis

Database search for polymorphisms

Genes within the reduced region of *Inf2* (34.38–35.34 Mbp) were searched for known exon (non-synonymous), intron (including splice sites) and untranslated region (UTR) polymorphisms using the public Mouse Phenome Database (table 1 in the online data supplement) [15]. *H2-Aa* and *H2-Ab1* genes are haplotype H2^K in the 2R and 4R congenic strains, thus excluding these as candidate genes [13]. Additionally, the *H2-Eb1* gene is divided between resistant and susceptible haplotypes and therefore remains a candidate gene, as well as *H2-Eb2* [16]. The genes that contain known non-synonymous polymorphisms between the B6 (susceptible) and C3 (resistant) mice in the reduced *Inf2* in order of location on chromosome 17 are *H2-Eb1*, *H2-Eb2*, *Btnl1*, *Notch4*, *Fkbp1*, *Crebl1*, *Tnxb*, *C4b*, *C4a*, *Stk19*, *Cfb*, *C2*, *Msh5*, *Ddah2*, *Ly6g6c*, *Bat5*, *Ly6g5c*, *Csnk2b*, *Aif1*, *Tnf* and *Lta* (table 1 in the online data




FIGURE 2. Bronchoalveolar lavage (BAL) fluid inflammatory parameters in the C3H-H2^b (□) and HeSnJ (■) mice exposed continuously to air or to 0.3 ppm O₃ for 48 or 72 h. a) Macrophages, b) polymorphonuclear leukocytes (PMNs), c) epithelial cells and d) total BAL fluid protein. Data are presented as mean ± SEM (n=3–5 per experimental group). *: significantly different from air-exposed mice (p<0.05); #: significantly different from HeSnJ mice (p<0.05).

supplement). UTR polymorphisms between the B6 and C3 mice were found in *H2-Eb2*, *Btnl2*, *Ppt2*, *C4b*, *C4a*, *G6b*, *Bat5*, *Ly6g5b*, and *Tnf* (table 1 in the online data supplement). Information is absent for *Lta* and *Tnf* UTR and intronic polymorphisms in the C3 strain, thus it is possible that more exist between the B6 and C3 strains.

We then asked if mRNA expression differences between B6 and C3 mice existed for several candidate genes within the reduced *Inf2*, and with either known or possible single nucleotide polymorphisms (fig. 4). *H2-Eb1* expression was reduced after 24 h of O₃ exposure in the C3 mice, but unchanged in the B6 mice; *H2-Eb2* expression was significantly increased only in B6 mice at 24 h. Expression of *Hspa1b* and *Btnl1* was significantly increased after 72 h O₃ in B6 mice only. *Tnf* and *Lta* expression was significantly increased in B6 mice after 24 h O₃ compared with no increases observed in C3 mice.

MHC class II deficient mice

Because of the prevalence of MHC class II genes in *Inf2* and the non-synonymous, intronic and UTR polymorphisms in *H2-Eb1* and *-Eb2* between the susceptible (B6) and resistant strains (C3) (table 1 in the online supplementary material), we hypothesised that they were candidate susceptibility genes for

O₃-induced inflammation. *H2Abl-Ea*^{-/-} mice are deficient in MHC class II genes *H2-Ab1*, *-Aa*, *-Eb1*, *-Eb2* and *-Ea* compared to the *H2Abl-Ea*^{+/+} mice. The B6 background of these mice is null at *H2-Ea*, therefore *H2Abl-Ea*^{-/-} mice only test the importance of the *H2-Ab1* through *H2-Eb2* MHC class II region. Relative to respective air controls, O₃ caused significant (p<0.05) increases in mean numbers of BALF PMNs and macrophages in *H2Abl-Ea*^{+/+} mice after 24, 48 and 72 h, and after 24 and 48 h O₃ in *H2Abl-Ea*^{-/-} mice (fig. 5a and b, and table 2). However, significantly greater numbers of PMNs and macrophages were found in *H2Abl-Ea*^{+/+} mice compared with *H2Abl-Ea*^{-/-} mice after 24–72 h and 48 and 72 h exposures, respectively. Significant strain and O₃ effects on BAL epithelial cells were found after 48 h O₃ (table 2). Significant strain differences in BAL protein content were also found after 48 h O₃ (fig. 5a). mRNA expression of *Tnf* and *Lta* was significantly increased in the *H2Abl-Ea*^{+/+} mice after 24 h O₃; however, no increase was observed in the *H2Abl-Ea*^{-/-} mice (fig. 5c and d).

Mice deficient in *Lta* or the *Tnf* cluster

Based on the existing evidence to support a role for the *Tnf* cluster in the O₃ response [10], as well as the known intronic and UTR polymorphisms in *Tnf* and *Lta* genes between the B6




FIGURE 3. Bronchoalveolar lavage (BAL) fluid inflammatory parameters in B10 (■), 4R (■), 5R (■), 2R (□) and A/Wy (■) mice exposed continuously to air or to 0.3 ppm O₃ for 48 or 72 h. a) Macrophages, b) polymorphonuclear leukocytes (PMNs), c) epithelial cells and d) total BAL fluid protein. Data are presented as mean \pm SEM (n=3–12 per experimental group). *: significantly different from air-exposed mice ($p < 0.05$); #: significantly different from B10 mice ($p < 0.05$).

and HeJ mice, we tested the role of the *Tnf* cluster in the model. *Lta*^{-/-}, *Lta/Tnf/Ltb*^{-/-}, and wild-type mice were exposed continuously to air or 0.3 ppm O₃ for 24, 48 or 72 h. PMNs were significantly elevated in the *Lta*^{+/+} and *Lta/Tnf/Ltb*^{+/+} mice compared with mice deficient in *Lta* or the *Lta/Tnf/Ltb* cluster following 24 and 48 h O₃ for both strains, and 72 h exposure for *Lta/Tnf/Ltb*^{-/-} mice compared with air controls ($p < 0.05$; fig. 6a–d). Mean numbers of BALF macrophages were also significantly elevated in wild-type mice compared with *Lta*^{-/-} and *Lta/Tnf/Ltb*^{-/-} mice following 24–72 h O₃ compared with air controls ($p < 0.05$; table 2). Epithelial cell numbers were not different between exposure or strains (table 2). The BAL protein content was not affected by these genes, except for 72 h exposure in the *Lta*^{-/-} mice, supporting the hypothesis that different genes regulate these specific phenotypes (fig. 6a–d) [17].

MHC class II gene expression was also significantly different between the strains deficient in the *Tnf* cluster. *H2-Eb1* mRNA was unchanged in B6 wild-type mice; however, this gene was significantly decreased after 48 h O₃ in the *Lta*^{-/-} and *Lta/Tnf/Ltb*^{-/-} mice compared with controls (fig. 6e and f). *H2-Eb2* mRNA was significantly increased in B6 mice after 24 and 48 h O₃, and was significantly greater after O₃ than mRNA expression in *Lta*^{-/-} and *Lta/Tnf/Ltb*^{-/-} mice at these time points.

DISCUSSION

The overall objective of this investigation was two-fold. We first sought to restrict the length of *Inf2* to elucidate candidate susceptibility genes for O₃-induced inflammation. The second objective was to validate the role(s) of identified gene candidates. Significant differences in O₃-induced inflammation between C3.SW-H2b/SJ and C3H/HeSnJ strains confirmed the role of *Inf2* inasmuch as the only difference between these two strains is a congenic region of chromosome 17 that encompasses *Inf2*. Comparison of inflammatory responses in 2R, 4R and 5R congenic mice with respective background strains further confirmed the importance of *Inf2*, and more importantly reduced *Inf2* to 0.96 Mbp. The reduced *Inf2* includes MHC class I, II, and III genes, and non-MHC genes, some of which have previously been identified as candidate genes in other lung injury models (e.g. *C4a* [18]). In the current study, gene expression and sequence analyses of the reduced *Inf2* suggested that MHC class II genes and the *Tnf* cluster may be important in O₃-induced inflammation, and significantly different phenotypes between O₃-exposed *H2Abl-Ea*^{+/+} and *H2Abl-Ea*^{-/-} mice and the mice deficient in *Lta* alone or the entire *Tnf* cluster confirmed a role for these genes. To our knowledge, these are the first studies to conclusively demonstrate a role for MHC class II genes and the entire *Tnf* cluster in




FIGURE 4. Gene expression for candidate genes identified within the reduced *Inf2* region of B6 (susceptible; ■) and C3 (HeJ, resistant; □) mice. Six genes were analysed using quantitative RT-PCR: a) *H2Eb1*, b) *Btn1*, c) *H2Eb2*, d) *Tnf*, e) *Hspa1b* and f) *Lta*. Data are presented as mean \pm SEM (n=3–7 per experimental group) and were determined using the comparative C_T method (Materials and methods section of the online supplementary material). The y-axis represents the gene of interest normalised first to 18S followed by determination of the fold-change relative to the average air control value for each strain. *: significantly different from air-exposed mice (p<0.05); #: significantly different from B6 mice (p<0.05).

oxidant-induced lung inflammation and provide evidence supporting a susceptibility “superlocus”.

Genetic association studies have implicated several gene categories, such as pro-inflammatory cytokine genes [9], metabolism genes [19] and innate immunity genes, in responses to environmental stimuli in human populations [20].

In inbred mouse models, positional cloning approaches identified QTLs for a number of lung diseases [21]. However, while identification of QTLs is an important initial step in understanding the genetic determinants of disease, QTLs can contain hundreds of genes. It is, therefore, necessary to refine the disease QTLs to a limited set of genes that can be evaluated [22]. In our integrative genomics approach to reducing *Inf2* and




FIGURE 5. Deficiency in major histocompatibility complex class II genes significantly reduced O₃-induced responses. a) Bronchoalveolar lavage (BAL) fluid polymorphonuclear leukocytes (PMNs) and b) total protein in *H2Ab1-Ea*^{-/-} (□) and *H2Ab1-Ea*^{+/+} (■) mice exposed continuously to air or to 0.3 ppm O₃ for 24, 48 or 72 h. Data are presented as mean ± SEM (n=3–10 per experimental group). Gene expression for c) *Lta* and d) *Tnf* genes in *H2Ab1-Ea* deficient mice (□) compared with wild-type mice (■). Data are presented as mean ± SEM (n=3–6 per experimental group) and were determined using the comparative C_T method (Materials and methods section of the online supplementary material). The y-axis represents either *Lta* or *Tnf* normalised first to 18S followed by determination of the fold-change relative to the average air control value for each strain. *: significantly different from air-exposed mice (p<0.05); #: significantly different from *H2Ab1-Ea*^{+/+} mice (p<0.05).

identifying candidate genes, we used congenic strains that were developed initially for histo-incompatibility studies by repeated backcrossing of regions of chromosome 17 from the donor strain onto the recipient background strain [23]. One advantage of these strains is that the allelic designations are well characterised and, after phenotyping these strains for their inflammatory response to O₃, it enabled restriction of the previously described *Inf2* QTL to a limited chromosomal interval to identify gene candidates.

We queried the Mouse Phenome Database to identify genes located within the narrowed *Inf2* that are polymorphic between susceptible B6 and resistant C3 mice. Several of the MHC class II and III genes (and those non-MHC genes located within these regions) have non-synonymous polymorphisms in exons, many have intronic polymorphisms, and several have UTR polymorphisms. However, not all of the genes have been assessed for polymorphisms between B6 and C3 mice (e.g. *Hspa1b*). It is important to note that functional polymorphisms may also exist in intronic regions, such as with *K-ras* and

mouse lung cancer [24]. Future studies will delineate the functionality of polymorphic regions between the B6 and C3 mice in those genes in which the functionality is unknown, such as *Ltb*.

Organisation of the H2 locus

Genes in the mouse *H2* locus and the homologous human *MHC* locus are organised in the same relative gene order [25] and span several Mbp. The *H2* locus contains class I, class II and class III genes. Class I and class II genes are important in fulfilling immunological functions. Class I molecules are expressed by most cells, and present endogenous antigens (cytosolic-derived) to CD8⁺ cytotoxic T-cells. Class II molecules (e.g. *H2-Eb2*, *Btl1*) are expressed in antigen presenting cells, such as dendritic cells, and present exogenous, endocytically derived antigens to CD4⁺ helper T-cells [26]. Class III genes are diverse in functionality, are located between the class I and class II genes and include the complement component genes (*C2* and *C4*), heat shock proteins (*Hspa1a*, *Hspa1b*, and *Hsc70t*) and the *Tnf* cluster.




FIGURE 6. O₃-induced responses in mice deficient in *Lta* or the *Tnf* cluster (*Lta*, *Tnf*, *Ltb*) genes. Bronchoalveolar lavage (BAL) fluid a) polymorphonuclear leukocytes (PMNs) and b) total protein for mice deficient in *Lta* (□) and their wild-type controls (■). Data are presented as mean ± SEM (n=3–10 per experimental group). *: significantly different from air-exposed mice (p<0.05); #: significantly different from *Lta*^{+/+} mice (p<0.05). c and d) BAL fluid inflammatory parameters for the *Lta/Tnf/Ltb* deficient mice (■) in response to O₃ compared to wild-type mice (□). Data are presented as mean ± SEM (n=3–10 per experimental group). *: significantly different from air-exposed mice (p<0.05); #: significantly different from *Lta/Tnf/Ltb*^{+/+} mice (p<0.05). Gene expression for the major histocompatibility complex class II genes e) *H2-Eb1* and f) *H2-Eb2* in the mice deficient in either *Lta* (■) or *Lta/Tnf/Ltb* (■) compared with the wild-type mice (■). B6 are the wild type for both strains and the numbers were combined since they were not significantly different from one another. No 24 h samples for *Lta/Tnf/Ltb*^{-/-} mice were done for gene expression analysis. Data are presented as means ± SEM (n=3–6 per experimental group) and were determined using the comparative C_T method (Materials and methods section of the online supplementary material). The y-axis represents either *H2-Eb1* or *H2-Eb2* normalised first to 18S followed by determination of the fold-change relative to the average air control value for each strain. *: significantly different from air-exposed mice (p<0.05); #: significantly different from wild-type mice (p<0.05).

TABLE 2 Effect of deletion of *H2-Abl-Ea*, *Lta* or *Lta/Tnf/Ltb* on macrophages and epithelial cells recovered from bronchoalveolar lavage (BAL) following O₃ exposure

Cell type	Genotype	Air [#]	O ₃ exposed		
			24 h	48 h	72 h
Macrophages	<i>H2Abl-Ea</i> ^{+/+}	19.6±0.1	33.6±3.9*	48.1±0.4* [¶]	37.9±3.7* [¶]
	<i>H2Abl-Ea</i> ^{-/-}	19.0±0.1	38.5±6.3*	33.2±0.4*	22.5±1.3
	<i>Lta</i> ^{+/+}	12.1±1.2	22.2±3.9* [¶]	29.7±7.7* [¶]	37.9±3.7*
	<i>Lta</i> ^{-/-}	13.8±1.9	7.6±2.2	21.2±2.7	32.0±2.5*
	<i>Lta/Tnf/Ltb</i> ^{+/+}	18.3±3.4	26.7±3.0	36.8±3.9* [¶]	38.5±3.4* [¶]
	<i>Lta/Tnf/Ltb</i> ^{-/-}	26.6±3.2	32.7±3.7	25.2±3.0	24.4±3.9
Epithelial cells	MHC ^{+/+}	1.4±0.3	1.0±0.2	2.6±0.7* [¶]	1.0±0.3
	MHC ^{-/-}	0.1±0.2	1.8±0.6	0.9±0.1	0.9±0.2
	<i>Lta</i> ^{+/+}	0.7±0.2	0.85±0.3	1.4±0.2	1.0±0.3
	<i>Lta</i> ^{-/-}	0.4±0.2	0.5±0.2	1.2±0.3	1.7±0.2
	<i>Lta/Tnf/Ltb</i> ^{+/+}	1.0±0.5	0.6±0.04	1.5±0.3 [¶]	1.8±1.5
	<i>Lta/Tnf/Ltb</i> ^{-/-}	1.8±1.5	1.0±0.3	0.6±0.05	1.7±0.3

Data are presented as mean±SEM number of cells × 10³ per mL BAL, n=3–10 mice per treatment group. [#]: air controls shown are for the 72-h time point except for the *Tnf*^{+/+} and *Tnf*^{-/-} mice (48-h time point shown). Air controls for each time point were not significantly different from one another. *: p<0.05, significantly elevated numbers of cells compared to air controls; [¶]: p<0.05, significantly elevated numbers of cells compared to the transgenic strain.

Candidate gene analysis

Deficiency in MHC class II genes and the *Tnf* cluster decreased the O₃-induced PMN and macrophage phenotypes, similar to those reductions seen in the 2R and 5R mice. Previous studies, as well as the data presented here, support a clear role for the *Tnf* cluster in these O₃-induced phenotypes. The *H2-Ea* gene (which is deficient in B6 mice), *H2-Ab1* and *H2-Aa* (as described earlier) can be excluded as candidate susceptibility genes.

CHEN *et al.* [27] demonstrated that inhibition of the CD4+ T-cell population using a monoclonal CD4 antibody *in vivo* greatly reduced the O₃-induced phenotypes (including PMNs, lymphocytes and epithelial cells) in B6 mice after 72 h continuous O₃ exposure. These studies support our findings in the *H2Abl-Ea*^{-/-} mice and suggest that, in O₃ susceptible strains, CD4+ T-cells are critical to O₃-induced inflammation and injury. A recent study in human airway monocytes isolated from exposed individuals suggests that O₃ primes the airway monocytes for innate immune responses, increases the capacity of the monocytes to present antigen to the CD4+ T-cells and increases the overall population of antigen presenting cells in the lungs [20].

Candidate susceptibility genes in *Inf2* and human disease

The MHC class II genes and the *Tnf* cluster in the reduced *Inf2* are all located in close proximity on chromosome 17 in mice and chromosome 6 in humans (fig. 1; table 1 of the online supplementary material). Several of these homologous genes have been implicated in human association studies. For example, polymorphisms in *HLA-DRB1* (*H2-Eb1*), *TNF* and *LTA* have been associated with sarcoidosis [28], a chronic granulomatous disease of unknown aetiology, supporting a role for a susceptibility "superlocus". In addition, *TNF* is a susceptibility gene for O₃-induced changes in lung function in humans [29]. Thus, it is possible that a cluster of genes in the

MHC class II region (*H2-Eb1* and *H2-Eb2*) interacts with the *Tnf* cluster (*Tnf*, *Lta* and *Ltb*) to promote O₃-induced lung inflammation and injury. In support of this notion, the MHC (specifically HLA genes) has the strongest influence on susceptibility to human autoimmune diseases, but recent evidence suggests that *TNF* may also be involved in autoimmune diseases [30].

In summary, this integrative genomics investigation utilised congenic mouse lines to narrow the chromosome 17 QTL for susceptibility to O₃-induced inflammation in the mouse. We used sequence analysis to identify candidate genes in the narrowed QTL, which were then tested for proof of concept. The novel role of MHC Class II genes and the *Tnf* cluster in susceptibility to O₃-induced inflammation provides unique insight to the mechanisms of O₃ effects in the lung and may lead to alternative means to prevent oxidant injury to lung tissues.

SUPPORT STATEMENT

This research was supported by the Intramural Research program of the National Institute of Environmental Health Sciences, National Institutes of Health, Dept of Health and Human Services (S.R. Kleeberger, H-Y. Cho, C. Walker, S. Tivedi, W. Gladwell) and Michigan State University (A.K. Bauer and E.A. Rondini).

STATEMENT OF INTEREST

A statement of interest for this study can be found at www.erj.ersjournals.com/misc/statements.dtl

ACKNOWLEDGEMENTS

Ozone exposures were conducted at the National Institute of Environmental Health Sciences Inhalation Facility under contract to Alion Science and Technology, Inc. (Research Triangle Park, NC, USA). The authors thank D. Morgan and H. Price at Alion Science and Technology, Inc. for coordinating the inhalation exposures. We also thank D. Cook (NIEHS, Research Triangle Park, NC, USA) for reviewing the manuscript and providing excellent suggestions.

REFERENCES

- 1 Delfino RJ, Murphy-Moulton AM, Burnett RT, *et al.* Effects of air pollution on emergency room visits for respiratory illnesses in Montreal, Quebec. *Am J Respir Critical Care Med* 1997; 155: 568–576.
- 2 Romieu I, Sienra-Monge JJ, Ramírez-Aguilar M, *et al.* Genetic polymorphism of GSTM1 and antioxidant supplementation influence lung function in relation to ozone exposure in asthmatic children in Mexico City. *Thorax* 2004; 59: 8–10.
- 3 US EPA. Air Quality Criteria for Ozone and Related Photochemical Oxidants (2006 Final). Washington, US Environmental Protection Agency, 2006. EPA/600/R-05/004aF-cF.
- 4 Goldstein BD, Lai LY, Ross SR, *et al.* Susceptibility of inbred mouse strains to ozone. *Arch Environ Health* 1973; 27: 412–413.
- 5 Kleeberger SR, Levitt RC, Zhang LY, *et al.* Linkage analysis of susceptibility to ozone-induced lung inflammation in inbred mice. *Nat Genet* 1997; 17: 475–478.
- 6 Prows DR, Shertzer HG, Daly MJ, *et al.* Genetic analysis of ozone-induced acute lung injury in sensitive and resistant strains of mice. *Nat Genet* 1997; 17: 471–474.
- 7 Alexis NE, Zhou H, Lay JC, *et al.* The glutathione-S-transferase Mu 1 null genotype modulates ozone-induced airway inflammation in human subjects. *J Allergy Clin Immunol* 2009; 124: 1222–1228.
- 8 Bergamaschi E, De Palma G, Mozzoni P, *et al.* Polymorphism of quinone-metabolizing enzymes and susceptibility to ozone-induced acute effects. *Am J Respir Crit Care Med* 2001; 163: 1426–1431.
- 9 Lee YL, McConnell R, Berhane K, *et al.* Ambient ozone modifies the effect of tumor necrosis factor G-308A on bronchitic symptoms among children with asthma. *Allergy* 2009; 64: 1342–1348.
- 10 Cho HY, Zhang LY, Kleeberger SR. Ozone-induced lung inflammation and hyperreactivity are mediated *via* tumor necrosis factor-alpha receptors. *Am J Physiol Lung Cell Mol Physiol* 2001; 280: L537–L546.
- 11 Stewart CA, Horton R, Allcock RJ, *et al.* Complete MHC haplotype sequencing for common disease gene mapping. *Genome Res* 2004; 14: 1176–1187.
- 12 Mouse Genome Informatics. www.informatics.jax.org/javawi2/wifetch?page=markerQF Date last accessed: July 21, 2009. Date last updated: May 19, 2010.
- 13 Laboratories J. H2 Haplotype Appendix 2. http://jaxmice.jax.org/literature/catalog/mhc_h2_haplotypes.pdf Date last accessed: March 2, 2009.
- 14 Bauer AK, Dixon D, DeGraff LM, *et al.* Toll-like receptor 4 in butylated hydroxytoluene-induced mouse pulmonary inflammation and tumorigenesis. *J Nat Cancer Inst* 2005; 97: 1778–1781.
- 15 Laboratory MPPatJ. Mouse Phenome Database. www.phenome.jax.org/SNP Date last accessed: July 1, 2009. Updated 2010.
- 16 Hood L, Steinmetz M, Malissen B. Genes of the major histocompatibility complex of the mouse. *Annu Rev Immunol* 1983; 1: 529–568.
- 17 Kleeberger SR, Reddy S, Zhang LY, *et al.* Genetic susceptibility to ozone-induced lung hyperpermeability: role of toll-like receptor 4. *Am J Respir Cell Mol Biol* 2000; 22: 620–627.
- 18 Haston CK, Tomko TG, Godin N, *et al.* Murine candidate bleomycin induced pulmonary fibrosis susceptibility genes identified by gene expression and sequence analysis of linkage regions. *J Med Genet* 2005; 42: 464–473.
- 19 Islam T, Berhane K, McConnell R, *et al.* Glutathione-S-transferase (GST) P1, GSTM1, exercise, ozone and asthma incidence in school children. *Thorax* 2009; 64: 197–202.
- 20 Lay JC, Alexis NE, Kleeberger SR, *et al.* Ozone enhances markers of innate immunity and antigen presentation on airway monocytes in healthy individuals. *J Allergy Clin Immunol* 2007; 120: 719–722.
- 21 Bauer AK, Malkinson AM, Kleeberger SR. Susceptibility to neoplastic and non-neoplastic pulmonary diseases in mice: genetic similarities. *Am J Physiol Lung Cell Mol Physiol* 2004; 287: L685–L703.
- 22 Kleeberger SR, Schwartz DA. From quantitative trait locus to gene: a work in progress. *Am J Respir Crit Care Med* 2005; 171: 804–805.
- 23 Silver L. Mouse Genetics. Concepts and Applications. New York, Oxford University Press, 1995.
- 24 Chen B, Johanson L, Wiest JS, *et al.* The second intron of the K-ras gene contains regulatory elements associated with mouse lung tumor susceptibility. *Proc Natl Acad Sci USA* 1994; 91: 1589–1593.
- 25 Snoek M, Groot PC, Spies T, *et al.* Fine mapping of the crossover-sites in the C4-H-2D region of H-2 recombinant mouse strains. *Immunogenetics* 1991; 34: 409–412.
- 26 Snoek M, Teuscher C, van Vugt H. Molecular analysis of the major MHC recombinational hot spot located within the G7c gene of the murine class III region that is involved in disease susceptibility. *J Immunol* 1998; 160: 266–272.
- 27 Chen X, Gavett SH, Wills-Karp M. CD4+ T lymphocyte modulation of ozone-induced murine pulmonary inflammation. *Am J Respir Cell Mol Biol* 1995; 12: 396–403.
- 28 Mrazek F, Holla LI, Hutyrova B, *et al.* Association of tumour necrosis factor-alpha, lymphotoxin-alpha and HLA-DRB1 gene polymorphisms with Lofgren's syndrome in Czech patients with sarcoidosis. *Tissue Antigens* 2005; 65: 163–171.
- 29 Yang IA, Holz O, Jorres RA, *et al.* Association of tumor necrosis factor-alpha polymorphisms and ozone-induced change in lung function. *Am J Respir Crit Care Med* 2005; 171: 171–176.
- 30 Camarena A, Juarez A, Mejia M, *et al.* Major histocompatibility complex and tumor necrosis factor-alpha polymorphisms in pigeon breeder's disease. *Am J Respir Crit Care Med* 2001; 163: 1528–1533.